







Cell 1 Regional Coastal Monitoring Programme Analytical Report 16: 'Full Measures' Survey 2023



North Tyneside Council
October 2023

Contents

Discl	aimer	i
Abbr	eviations and Acronyms	ii
	er Levels Used in Interpretation of Changes	
	sary of Terms	
Prea	mble	iv
1. I	Introduction	9
1.1	Study Area	9
1.2	Methodology	9
2. /	Analysis of Survey Data	10
2.1	Whitley Sands	
2.2	Cullercoats Bay	12
2.3	Tynemouth Long Sands	
2.4	King Edward's Bay	
3. I	Problems Encountered and Uncertainty in Analysis	
	Recommendations for 'Fine-tuning' the Monitoring Programme	
	Conclusions and Areas of Concern	

AppendicesAppendix A
Appendix B **Beach Profiles** Topographic Survey

List of Figures

Figure 1 Sediment Cells in England and Wales

Figure 2 **Survey Locations**

List of Tables

Analytical, Update and Overview Reports Produced to Date Sub-division of the Cell 1 Coastline Table 1

Table 2

Authors	
Alix Scullion	Royal HaskoningDHV
Dr Nick Cooper – Review & Approval	Royal HaskoningDHV

Disclaimer

Royal HaskoningDHV has prepared this report in accordance with the instructions of our client Scarborough Borough Council (SBC) for the client's sole and specific use. Any other persons who use any information contained herein do so at their own risk. Royal HaskoningDHV has used reasonable skill, care and diligence in the interpretation of data provided to them and accepts no responsibility for the content, quality or accuracy of any Third party reports, monitoring data or further information provided either to them by SBC or, via SBC from a Third party source, for analysis under this term contract.

Data and reports collected as part of the Cell 1 Regional Coastal Monitoring Programme are available to download via the North East Coastal Observatory via the webpage: www.northeastcoastalobservatory.org.uk.

The North East Coastal Observatory does not "license" the use of images or data or sign license agreements. The North East Coastal Observatory generally has no objection to the reproduction and use of these materials (aerial photography, wave data, beach surveys, bathymetric surveys, reports), subject to the following conditions:

- 1. North East Coastal Observatory material may not be used to state or imply the endorsement by North East Coastal Observatory or by any North East Coastal Observatory employee of a commercial product, service, or activity, or used in any manner that might mislead.
- 2. North East Coastal Observatory should be acknowledged as the source of the material in any use of images and data accessed through this website, please state "Image/Data courtesy of North East Coastal Observatory". We recommend that the caption for any image and data published includes our website, so that others can locate or obtain copies when needed. We always appreciate notification of beneficial uses of images and data within your applications. This will help us continue to maintain these freely available services. Send e-mail to Robin.Siddle@scarborough.gov.uk
- 3. It is unlawful to falsely claim copyright or other rights in North East Coastal Observatory material.
- 4. North East Coastal Observatory shall in no way be liable for any costs, expenses, claims, or demands arising out of the use of North East Coastal Observatory material by a recipient or a recipient's distributees.
- 5. North East Coastal Observatory does not indemnify nor hold harmless users of North East Coastal Observatory material, nor release such users from copyright infringement, nor grant exclusive use rights with respect to North East Coastal Observatory material.
- 6. North East Coastal Observatory material is not protected by copyright unless noted (in associated metadata). If copyrighted, permission should be obtained from the copyright owner prior to use. If not copyrighted, North East Coastal Observatory material may be reproduced and distributed without further permission from North East Coastal Observatory.

Abbreviations and Acronyms

Acronym / Abbreviation	Definition	
AONB	Area of Outstanding Natural Beauty	
DGM	Digital Ground Model	
HAT	Highest Astronomical Tide	
LAT	Lowest Astronomical Tide	
MHWN	Mean High Water Neap	
MHWS	Mean High Water spring	
MLWS	Mean Low Water Neap	
MLWS	Mean Low Water spring	
m	metres	
ODN	Ordnance Datum Newlyn	

Water Levels Used in Interpretation of Changes

	Water Level (m AOD)
Water Level Parameter	Whitley Sands to
	King Edward's Bay
HAT	3.1
MHWS	2.4
MHWN	1.3
MLWN	-0.8
MLWS	-1.9

Source: UKHO Admiralty Tide Tables, 2020

Glossary of Terms

Term	Definition	
Beach nourishment	Artificial process of replenishing a beach with material from another source.	
Berm crest	Ridge of sand or gravel deposited by wave action on the shore just above the normal high water mark.	
Breaker zone	Area in the sea where the waves break.	
Coastal squeeze	The reduction in habitat area which can arise if the natural landward migration of a habitat under sea level rise is prevented by the fixing of the high water mark, e.g. a sea wall.	
Downdrift	Direction of alongshore movement of beach materials.	
Ebb-tide	The falling tide, part of the tidal cycle between high water and the next low water.	
Fetch	Length of water over which a given wind has blown that determines the size of the waves produced.	
Flood-tide	Rising tide, part of the tidal cycle between low water and the next high water.	
Foreshore	Zone between the high water and low water marks, also known as the intertidal zone.	
Geomorphology	The branch of physical geography/geology which deals with the form of the Earth, the general configuration of its surface, the distribution of the land, water, etc.	
Groyne	Shore protection structure built perpendicular to the shore; designed to trap sediment.	
Mean High Water (MHW)	The average of all high waters observed over a sufficiently long period.	
Mean Low Water (MLW)	The average of all low waters observed over a sufficiently long period.	
Mean Sea Level (MSL)	Average height of the sea surface over a 19-year period.	
Offshore zone	Extends from the low water mark to a water depth of about 15 m and is permanently covered with water.	
Storm surge	A rise in the sea surface on an open coast, resulting from a storm.	
Swell	Waves that have travelled out of the area in which they were generated.	
Tidal prism	The volume of water within the estuary between the level of high and low tide, typically taken for mean spring tides.	
Tide	Periodic rising and falling of large bodies of water resulting from the gravitational attraction of the moon and sun acting on the rotating earth.	
Topography	Configuration of a surface including its relief and the position of its natural and man-made features.	
Transgression	The landward movement of the shoreline in response to a rise in relative sea level.	
Updrift	Direction opposite to the predominant movement of longshore transport.	
Wave direction	Direction from which a wave approaches.	
Wave refraction	Process by which the direction of approach of a wave changes as it moves into shallow water.	

Preamble

The Cell 1 Regional Coastal Monitoring Programme covers approximately 300km of the north east coastline, from the Scottish Border (just south of St. Abb's Head) to Flamborough Head in East Yorkshire. This coastline is often referred to as 'Coastal Sediment Cell 1' in England and Wales (Figure 1). Within this frontage the coastal landforms vary considerably, comprising low-lying tidal flats with fringing salt marshes, hard rock cliffs that are mantled with glacial sediment to varying thicknesses, softer rock cliffs and extensive landslide complexes.




Figure 1 Sediment Cells in England and Wales

The programme commenced in its present guise in September 2008¹ and is managed by Scarborough Borough Council on behalf of the North East Coastal Observatory. It is funded by the Environment Agency, working in partnership with the following organisations:



Royal HaskoningDHV has been appointed to provide Analytical Services in relation to the present phase of the Cell 1 Regional Coastal Monitoring Programme, between 2016 - 2027.

The main elements of the Cell 1 Regional Coastal Monitoring Programme involve:

¹ Prior to 2008, coastal monitoring was undertaken on a consistent basis across Northumberland and North Tyneside as part of the (then) Northumbrian Coastal Authorities Group's monitoring programme which commenced in 2002, whilst several authorities between the River Tyne and Flamborough Head undertook their own local monitoring programmes.

- beach profile surveys
- topographic surveys
- cliff top recession surveys
- real-time wave data collection
- bathymetric and seabed characterisation surveys
- aerial photography
- LiDAR surveys
- walk-over cliff and coastal defence asset surveys

The beach profile surveys, topographic surveys and cliff top recession surveys are undertaken as a 'Full Measures' survey in autumn/early winter every year. Some of these surveys are then repeated the following spring as part of a 'Partial Measures' survey.

Each year, an Analytical Report is produced for each individual authority, providing a detailed analysis and interpretation of the 'Full Measures' surveys. This is followed by a brief Update Report for each individual authority, providing ongoing findings from the 'Partial Measures' surveys.

Annually, a Cell 1 Overview Report is also produced. This provides a region-wide summary of the main findings relating to trends and interactions along the entire Cell 1 frontage.

To date the following reports have been produced:

Table 1 Analytical, Update and Overview Reports Produced to Date

		Full Measures		Partial Measures		Cell 1
	Year	Survey	Analytical Report	Survey	Update Report	Overview Report
1	2008/09	Sep-Dec 08	May 09	Mar-May 09	Jun 09	
2	2009/10	Sep-Dec 09	Mar 10	Feb-Mar 10	Jul 10	
3	2010/11	Aug-Nov 10	Feb 11	Feb-Apr 11	Aug 11	Sep 11
4	2011/12	Oct-Nov 11	Oct 12	Mar-May 12	Feb 13	
5	2012/13	Sep-Oct 12	Mar 13	Mar-Apr 13	Jun 13	
6	2013/14	Sep-Oct 13	Feb 14	Mar-Apr 14	Jul 14	
7	2014/15	Oct-Nov 14	Feb 15	Mar 15	Jul 15	
8	2015/16	Oct-Nov 15	Feb 16	Mar 16	Jul 16	Jun 16
9	2016/17	Sep 16	Feb 17	Mar 17	Jul 17	
10	2017/18	Sep-Oct 17	Feb 18	Mar 18	May 18	
11	2018/19	Sep-Oct 18	Jan 19	Feb-Mar 19	Aug 19	
12	2019/20	Sep-Nov 19	Dec 19	Mar 20	Mar 20	
13	2020/21	Sep 20	Nov 20	Mar 21	Apr 21	Aug 21
14	2021/22	Aug 21	Dec 21	Mar-Apr 22	May 22	
15	2022/23	Aug 22	Jan 23	Feb-Mar 23	May 23	
16	2023/24	Aug 23(*)				

^(*) The present report is **Analytical Report 16** and provides an analysis of the 2023 Full Measures survey for North Tyneside Council's frontage.

In addition, separate reports are produced for other elements of the programme as and when specific components are undertaken, such as wave data collection, bathymetric and seabed sediment data collection, aerial photography, and walk-over visual inspections.

For purposes of analysis, the Cell 1 frontage has been split into the sub-sections listed in the Table 2.

Table 2 Sub-divisions of the Cell 1 Coastline

Authority	Zone
	Spittal A
	Spittal B
	Goswick Sands
	Holy Island
	Bamburgh
	Beadnell Village
Northumberlan	Beadnell Bay
d County	Embelton Bay
Council	Boulmer
	Alnmouth Bay
	High Hauxley and Druridge Bay
	Lynemouth Bay
	Newbiggin Bay
	Cambois Bay
	Blyth South Beach
North	Whitley Sands
Tyneside	Cullercoats Bay
Council —	Tynemouth Long Sands
Countri	King Edward's Bay
Courth	Littehaven Beach
South Tyneside	Herd Sands
Council —	Trow Quarry (incl. Frenchman's Bay)
Council	Marsden Bay
	Whitburn Bay
Sunderland	Harbour and Docks
Council	Hendon to Ryhope (incl. Halliwell Banks)
	Featherbed Rocks
Durham	Seaham
County	Blast Beach
Council	Hawthorn Hive
	Blackhall Colliery
Hartlepool	North Sands
Borough	Headland
Council	Middleton
000.1011	Hartlepool Bay
	Coatham Sands
Redcar &	Redcar Sands
Cleveland	Marske Sands
Borough	Saltburn Sands
Council	Cattersty Sands (Skinningrove)
	Staithes
	Staithes
	Runswick Bay
North	Sandsend Beach, Upgang Beach and Whitby Sands
Yorkshire	Robin Hood's Bay
Council —	Scarborough North Bay
	Scarborough South Bay
	Cayton Bay
	Filey Bay

1. Introduction

1.1 Study Area

North Tyneside Council's frontage extends from Hartley (just south of Blyth) in the north, to the River Tyne in the south. For the purposes of this report and for consistency with previous reporting, it has been sub-divided into four areas, namely:

- Whitley Sands
- Cullercoats Bay
- Tynemouth Long Sands
- King Edward's Bay

1.2 Methodology

Along North Tyneside Council's frontage, the following surveying is undertaken:

- Full Measures survey annually each autumn/early winter comprising:
 - Beach profile surveys along eight transect lines (commenced 2002)
 - Beach profile surveys along an additional two transects (commenced 2010)
 - o Topographic survey along Whitley Sands (commenced 2010)
 - o Topographic survey along Tynemouth Long Sands (commenced 2011)
- Partial Measures survey annually each spring comprising:
 - o Beach profile surveys along all ten transect lines (commenced 2010)

The location of these surveys is shown in Figure 1. The beach profiles and topographic surveys were undertaken between the 15th and 16th August 2023. The weather conditions varied throughout the survey; further details can be found in the surveyor's reports.


All data have been captured in a manner commensurate with the principles of the Environment Agency's *National Standard Contract and Specification for Surveying Services* and stored in a file format compatible with the software systems being used for the data analysis, namely SANDS and ArcGIS. This data collection approach and file format is comparable to that being used on other regional coastal monitoring programmes, such as in the South East and South West of England.

Upon receipt of the data from the survey team, they are quality assured and then uploaded onto the programme's website for storage and availability to others and also input to SANDS and GIS for subsequent analysis.

The Analytical Report is then produced following a standard structure for each authority. This involves:

- description of the changes observed since the previous survey and an interpretation of the drivers of these changes (Section 2);
- documentation of any problems encountered during surveying or uncertainties inherent in the analysis (Section 3);
- recommendations for 'fine-tuning' the programme to enhance its outputs (Section 4); and
- providing key conclusions and highlighting any areas of concern (Section 5).

Data from the present survey are presented in a processed form in the Appendices.



2. Analysis of Survey Data

2.1 Whitley Sands

Survey Date	Description of Changes Since Last Survey	Interpretation
	Beach Profiles: Whitley Sands is covered by five beach profile lines for the Full Measures survey (Appendix A). Four of these (1aNTDC01 to 1aNTDC04) were initially surveyed in April 2002 and were surveyed annually to 2009 (Full Measures, autumn 2009) and bi-annually thereafter. From March 2010 (Partial Measures, spring 2010) onwards, an additional beach profile line (NTDC04A) has been surveyed at the southern end of the frontage. All profiles were last surveyed in March 2023 for the Partial	Since the last survey, there has generally been accretion across the beach profiles except at Profile 1aNTDC04 which has lowered on the lower beach. The general trend indicates a landward movement of material, which is characteristic of summertime conditions.
	Measures survey.	Longer term trends: Beach levels are generally at a medium-high range compared to earlier surveys,
15 th August 2023	1aNTDC01 is located in the north of Whitley Sands, along the undefended cliffs immediately south of Trinity Road car park. The cliff was not measured due to dangerous access. The upper to middle beach has risen by up to 0.3m between chainages 28m to the rock patch at chainage 60m. Seaward of this point, there has been minor changes in beach profile, with rocks visible at chainages 92m and 101m. Overall, the toe of the cliff is at one of its most landward positions recorded. The upper beach is at a low level compared to the range recorded from previous surveys.	except profile 1aNTDC01 which is at a low level.
	Profile 1aNTDC02 is located in the northern part of Whitley Sands. A seawall is present at the back of the beach. The upper beach between the seawall at chainage 50m and chainage 71m has lowered by up to 0.2m. The majority of the beach profile seaward of this has risen by up to 0.5m to chainage 140m. The beach profile at the rock patch remains in a similar position to the previous survey. The profile is generally at a medium level compared with the range recorded from previous surveys.	
	Profile 1aNTDC03 is located at the centre of Whitley Sands. The beach profile has risen by up to 0.5m on the upper-middle beach, 0.3m on the middle beach and 0.1m on the lower beach. There is a small section of erosion between chainages 105-119m which has lowered by less than 0.1m. The profile is at a medium-high level compared to the range recorded from previous surveys.	
	Profile 1aNTDC04 is located in the southern part of Whitley Sands. There has been relatively little change between the seawall and chainage 33m. The beach profile between chainages 33-105m has	

Survey Date	Description of Changes Since Last Survey	Interpretation
	risen by up to 0.2m. Seaward of this point, the lower beach seaward of chainage 105m has lowered by up to 0.6m. The rock patch previously exposed at chainage 65m remains covered. Overall, the beach is at a high level compared with the range recorded in previous surveys.	
	Profile 1aNTDC04A is located towards the southern end of Whitley Sands. There has been accretion across the majority of the beach profile, by up to 0.7m on the upper beach and less than 0.1m on the middle and lower beach. A small section on the lower beach between chainages 97-124m has lowered by up to 0.1m. Overall, the profile is at a high level compared to the range recorded from previous surveys, particularly between chainage 70-97m which is at its highest level recorded.	
	Topographic Survey: Whitley Sands is covered by an annual topographic survey, which commenced in October 2010. Data from the most recent topographic survey (Full Measures, autumn 2023) have been used to create a digital ground model (DGM) (Appendix B – Map 1) using a GIS. A difference plot has also been produced using the DGM (Appendix B – Map 3) produced from the last produced topographic survey (Full Measures, autumn 2022) and the present survey.	The most recent topographic survey difference plot shows a mixture of erosion and accretion, however the most dominant trend is erosion on the upper to middle beach and accretion on the lower beach.
August 2023	The difference plot shows that the north of the beach is dominated by erosion on the upper-middle beach and little change on the lower beach. In the centre of the bay, the beach is similar ro the north with erosion on the upper to middle beach and accretion on the lower beach. The magnitude of change is greater in the centre of the bay, with erosion/accretion reaching up to ±1.5m. The beach in the south of the bay shows relatively little change on the upper to middle beach, with minor patches of erosion limited to 0.25m. The lower beach is dominated by accretion up to 1.0m.11xtent. Areas of accretion and erosion occur in shore-parallel bands. This is a comparison of annual surveys, rather than a biannual comparison like the beach profiles and therefore explains why the patterns do not match with those described above, which generally show accretion across the majority of the beach profiles.	

2.2 Cullercoats Bay

Survey Date	Description of Changes Since Last Survey	Interpretation
16 th August	Beach Profiles: Cullercoats Bay is covered by one beach profile line for the Full Measures survey (Appendix A). This was surveyed annually each autumn between 2002 and 2009. From spring 2010 onwards, it has been surveyed bi-annually.	As in previous surveys, access to the cliff has not been possible. The data shows only limited change related to short-term patterns of sediment movement.
2023	At profile 1aNTDC05 , the survey report notes that the cliff was not measured due to dangerous access. The beach profile has lowered on the upper beach between chainages 24-49m by up to 0.4m and on the lower beach between chainages 108-126m by up to 0.2m. The middle beach has risen by up to 0.3m. The profile is general at a low level on the upper beach, rising to a medium-high level on the middle and lower beach compared to the range recorded from previous surveys.	Longer term trends: The amount of change is within the past range. Much of the profile is at medium-high level compared with previous surveys.

2.3 Tynemouth Long Sands

Survey Date	Description of Changes Since Last Survey	Interpretation
15-16 th August 2023	Beach Profiles: Tynemouth Long Sands is covered by three beach profile lines for the Full Measures survey (Appendix A). Profiles 1aNTDC06 and 1aNTDC07 were initially surveyed annually each autumn between 2002 and 2009. A third profile, 1aNTDC06A, was then added in the centre of the frontage. From spring 2010 (Partial Measures) onwards, all profiles have been surveyed bi-annually. Profile 1aNTDC06 is located approximately 150m south of the access ramp towards the north of the bay. The top of the cliff has not changed since the last survey however, the remainder of the cliff profile cannot be reviewed due to lack of data points in the profile plot as the survey report notes 'no access to middle of section 6 due to seed protection fences'. The beach profile starts at c.35m chainage and shows an accumulation of sediment at the toe by up to 0.4m to chainage 48m. The upper beach between chainages 48-120m has lowered by up to 0.8m and up to 0.4m on the lower beach seaward of chainage 183m. The middle beach between chainages 120-183m has risen by up to 0.4mOverall, the profile is at a medium-high level compared with the range from previous surveys. At profile 1aNTDC06A, the dune-cliff face has not significantly changed in form or position. As with section 6, the survey report notes 'no access to middle of section 6a due to seed protection fences'. The upper beach between the dune toe and chainage 145m has remained stable since the previous survey. The rest of the beach profile has alternated between accretion by up to 0.5m and erosion by up to 0.2m. Overall, the autumn 2023 profile is at a high level compared to the range recorded from the previous surveys. Profile 1aNTDC07 is located approximately 50m south of the access route through the dunes towards the southern end of the bay. The dune-cliff has not significantly changed in form or position since the last survey. As with section 6 and 6a, the survey report notes 'no access to middle of section 7 due to seed protection fences'. There has been variable erosion an	Since the last survey the dunes have retained the same form and position. The profiles across Long Sands show variable erosion and accretion, associated with the seasonal movement of material. Longer term trends: Overall, the beaches have retained a similar form and are in the medium-high range of previous surveys.
August 2023	Topographic Survey:	The pattern shown in the difference plot supports the patterns identified in the profiles, with shore parallel

Survey Date	Description of Changes Since Last Survey	Interpretation
	The first survey was undertaken for the Full Measures survey in October 2010. Data from the current topographic survey have been used to create a digital ground model (DGM) (Appendix B – Map 2) using a Geographical Information System (GIS). A difference plot has also been produced by comparing the current DGM (Appendix B – Map 4) with that produced from the last topographic survey in February2023. The centre and north of the bay is dominated by shore parallel bands of erosion (on the mid-upper beach) and accretion (upper and mid-lower beach) with areas of little change (±0.1m) in-between. The south of the bay is dominated by low-level little change (±0.1m), with isolated patches of erosion and accretion. The magnitude of change is relatively small for both accretion and erosion in the south of the bay (±0.75m) but is of larger magnitude in the central and northern part of the bay (±1.75m).	bands of erosion and accretion, exhibiting berm movement.

2.4 King Edward's Bay

Survey Date	Description of Changes Since Last Survey	Interpretation
16 th August 2023	Beach Profiles: King Edward's Bay is monitored by one beach profile line for the Full Measures survey (Appendix A). This was surveyed annually between 2002 and 2009. From spring 2010 onwards, it has been surveyed bi-annually. At profile 1aNTDC08, the beach has generally risen across the profile. The beach between chainages 6-30m has risen by less than 0.1m, before switching to erosion between chainages 30-66m by up to 0.3m. The beach profile seaward of chainage 66m has risen by up to 0.45m to chainage 160m. The survey continues a further 60m, showing a berm on the lower beach at chainage 196m. Overall, the beach profile is at a medium-high level on the upper and middle beach and a low level on the lower beach when compared with the range recorded from previous surveys. The berm at the end of the survey is at a high level.	Since the last survey, the upper beach at King Edward's Bay has experienced movement of sediment in a seaward direction in the form of a berm. Longer term trends: The profile is at a medium-high level on the upper and middle beach, and a low level on the lower beach compared with the range of previously observed surveys at this location. Changes between the spring 2023 and autumn 2023 survey show a seasonal movement of sediment across the beach in the form of a berm.

3. Problems Encountered and Uncertainty in Analysis

Individual Profiles

- At profile 1aNTDC01 the cliff was not measured due to access problems.
- At profile 1aNTDC05 the cliff was not measured due to access problems. Access to this
 profile is noted to have been dangerous in previous Partial Measures and Full Measures
 reports, and it is recommended that the beach profile should start at the cliff toe and that
 the cliff be monitored using the aerial survey data.
- At Tynemouth Long Sands (profiles 1aNTDC06, 1aNTDC06A and 1aNTDC07) there was
 no access to the dunes in the middle of the profile due to seed protection fences. This
 means it has not yet been possible to monitor the effectiveness of the dune stabilisation
 scheme.

4. Recommendations for 'Fine-tuning' the Monitoring Programme

• It is recommended that access to the stabilised dunes at Tynemouth Long Sands be attempted in future surveys to monitor the effectiveness of the stabilisation fences.

5. Conclusions and Areas of Concern


- At Whitley Sands, there has been variable change across the beach profiles since the previous survey. For the most part, the beach is at a medium-high level in the range recorded from previous surveys. The topographic survey shows sediment movement from the upper and middle beach towards the lower beach.
- At Cullercoats Bay, there has been variable change across the beach profile, limited to ±0.4m. The results fall within the range of previously recorded results and there are no causes for concern.
- At Tynemouth Long Sands, the majority of recorded profiles are within the previously recorded range with berm building processes dominating. The profiles present no cause for concern. The topographic survey results generally demonstrate shore-parallel bands of erosion and accretion associated with seasonal distribution of sediment. Change in the centre and north of the bay is of larger magnitude (±1.75m) than the south of the bay (±0.75m).
- At King Edward's Bay, there appears to have been a seaward movement of sediment in the form of a berm. There are no causes for concern and the results fall within the range recorded in previous surveys.


Appendices


Appendix A Beach Profiles


The following sediment feature codes are used on some profile plots:


Code	Description
S	Sand
M	Mud
G	Gravel
GS	Gravel & Sand
MS	Mud & Sand
В	Boulders
R	Rock
SD	Sea Defence
SM	Saltmarsh
W	Water Body
GM	Gravel & Mud
GR	Grass
D	Dune (non-vegetated)
DV	Dune (vegetated)
F	Forested
X	Mixture
FB	Obstruction
CT	Cliff Top
CE	Cliff Edge
CF	Cliff Face
SH	Shell
ZZ	Unknown























































Appendix B Topographic Survey

