



**Cell 1 Regional Coastal Monitoring Programme
Analytical Report 9: 'Full Measures' Survey 2016**



Durham County Council

March 2017

Contents

Disclaimer	i
Abbreviations and Acronyms.....	ii
Water Levels Used in Interpretation of Changes	ii
Glossary of Terms.....	iii
Preamble	iv
1. Introduction	1
1.1 Study Area	1
1.2 Methodology.....	1
2. Analysis of Survey Data	1
2.1 Featherbed Rocks	1
2.2 Seaham (Dawdon)	2
2.3 Blast Beach.....	3
2.4 Hawthorne Hive.....	5
2.5 Blackhall Colliery	6
3. Problems Encountered and Uncertainty in Analysis	7
4. Recommendations for 'Fine-tuning' the Monitoring Programme	7
5. Conclusions and Areas of Concern	7

Appendices

Appendix A	Beach Profiles
Appendix B	Cliff Top Survey

List of Figures

Figure 1	Sediment Cells in England and Wales
Figure 2	Survey Location Maps

List of Tables

Table 1	Analytical, Update and Overview Reports Produced to Date
Table 2	Sub-division of the Cell 1 Coastline

Authors	
Emma Hick	Royal HaskoningDHV
Dr Nick Cooper – Review of Draft	Royal HaskoningDHV
Dr Nick Cooper – Approval of Final	Royal HaskoningDHV

Disclaimer

Royal HaskoningDHV has prepared this report in accordance with the instructions of our client Scarborough Borough Council (SBC) for the client's sole and specific use. Any other persons who use any information contained herein do so at their own risk. Royal HaskoningDHV has used reasonable skill, care and diligence in the interpretation of data provided to them and accepts no responsibility for the content, quality or accuracy of any Third party reports, monitoring data or further information provided either to them by SBC or, via SBC from a Third party source, for analysis under this term contract.

Data and reports collected as part of the Cell 1 Regional Coastal Monitoring Programme are available to download via the North East Coastal Observatory via the webpage: www.northeastcoastalobservatory.org.uk.

The North East Coastal Observatory does not "license" the use of images or data or sign license agreements. The North East Coastal Observatory generally has no objection to the reproduction and use of these materials (aerial photography, wave data, beach surveys, bathymetric surveys, reports), subject to the following conditions:

1. North East Coastal Observatory material may not be used to state or imply the endorsement by North East Coastal Observatory or by any North East Coastal Observatory employee of a commercial product, service, or activity, or used in any manner that might mislead.
2. North East Coastal Observatory should be acknowledged as the source of the material in any use of images and data accessed through this website, please state "Image/Data courtesy of North East Coastal Observatory". We recommend that the caption for any image and data published includes our website, so that others can locate or obtain copies when needed. We always appreciate notification of beneficial uses of images and data within your applications. This will help us continue to maintain these freely available services. Send e-mail to Robin.Siddle@scarborough.gov.uk
3. It is unlawful to falsely claim copyright or other rights in North East Coastal Observatory material.
4. North East Coastal Observatory shall in no way be liable for any costs, expenses, claims, or demands arising out of the use of North East Coastal Observatory material by a recipient or a recipient's distributees.
5. North East Coastal Observatory does not indemnify nor hold harmless users of North East Coastal Observatory material, nor release such users from copyright infringement, nor grant exclusive use rights with respect to North East Coastal Observatory material.
6. North East Coastal Observatory material is not protected by copyright unless noted (in associated metadata). If copyrighted, permission should be obtained from the copyright owner prior to use. If not copyrighted, North East Coastal Observatory material may be reproduced and distributed without further permission from North East Coastal Observatory.

Abbreviations and Acronyms

Acronym / Abbreviation	Definition
AONB	Area of Outstanding Natural Beauty
DGM	Digital Ground Model
HAT	Highest Astronomical Tide
LAT	Lowest Astronomical Tide
MHWN	Mean High Water Neap
MHWS	Mean High Water Spring
MLWS	Mean Low Water Neap
MLWS	Mean Low Water Spring
m	metres
ODN	Ordnance Datum Newlyn

Water Levels Used in Interpretation of Changes

Water Level Parameter	Water Level (m AOD)			
	River Tyne to Frenchman's Bay	Frenchman's Bay to Souter Point	Souter Point to Chourdon Point	Chourdon Point to Hartlepool Headland
1 in 200 year	3.41	3.44	3.66	3.91
HAT	2.85	2.88	3.18	3.30
MHWS	2.15	2.18	2.48	2.70
MLWS	-2.15	-2.12	-1.92	-1.90

Source: *River Tyne to Flamborough Head Shoreline Management Plan 2.*
Royal Haskoning, February 2007.

Glossary of Terms

Term	Definition
Beach nourishment	Artificial process of replenishing a beach with material from another source.
Berm crest	Ridge of sand or gravel deposited by wave action on the shore just above the normal high water mark.
Breaker zone	Area in the sea where the waves break.
Coastal squeeze	The reduction in habitat area which can arise if the natural landward migration of a habitat under sea level rise is prevented by the fixing of the high water mark, e.g. a sea wall.
Downdrift	Direction of alongshore movement of beach materials.
Ebb-tide	The falling tide, part of the tidal cycle between high water and the next low water.
Fetch	Length of water over which a given wind has blown that determines the size of the waves produced.
Flood-tide	Rising tide, part of the tidal cycle between low water and the next high water.
Foreshore	Zone between the high water and low water marks, also known as the intertidal zone.
Geomorphology	The branch of physical geography/geology which deals with the form of the Earth, the general configuration of its surface, the distribution of the land, water, etc.
Groyne	Shore protection structure built perpendicular to the shore; designed to trap sediment.
Mean High Water (MHW)	The average of all high waters observed over a sufficiently long period.
Mean Low Water (MLW)	The average of all low waters observed over a sufficiently long period.
Mean Sea Level (MSL)	Average height of the sea surface over a 19-year period.
Offshore zone	Extends from the low water mark to a water depth of about 15 m and is permanently covered with water.
Storm surge	A rise in the sea surface on an open coast, resulting from a storm.
Swell	Waves that have travelled out of the area in which they were generated.
Tidal prism	The volume of water within the estuary between the level of high and low tide, typically taken for mean spring tides.
Tide	Periodic rising and falling of large bodies of water resulting from the gravitational attraction of the moon and sun acting on the rotating earth.
Topography	Configuration of a surface including its relief and the position of its natural and man-made features.
Transgression	The landward movement of the shoreline in response to a rise in relative sea level.
Updrift	Direction opposite to the predominant movement of longshore transport.
Wave direction	Direction from which a wave approaches.
Wave refraction	Process by which the direction of approach of a wave changes as it moves into shallow water.

Preamble

The Cell 1 Regional Coastal Monitoring Programme covers approximately 300km of the north east coastline, from the Scottish Border (just south of St. Abb's Head) to Flamborough Head in East Yorkshire. This coastline is often referred to as 'Coastal Sediment Cell 1' in England and Wales (Figure 1). Within this frontage the coastal landforms vary considerably, comprising low-lying tidal flats with fringing salt marshes, hard rock cliffs that are mantled with glacial sediment to varying thicknesses, softer rock cliffs and extensive landslide complexes.




Figure 1 Sediment Cells in England and Wales

The work commenced with a three-year monitoring programme in September 2008 that was managed by Scarborough Borough Council on behalf of the North East Coastal Group. This initial phase has been followed by a five-year programme of work, which started in October 2011. The work is funded by the Environment Agency, working in partnership with the following organisations:



The main elements of the Cell 1 Regional Coastal Monitoring Programme involve:

- beach profile surveys
- topographic surveys
- cliff top recession surveys
- real-time wave data collection
- bathymetric and sea bed characterisation surveys
- aerial photography
- walk-over surveys

The beach profile surveys, topographic surveys and cliff top recession surveys are undertaken as a 'Full Measures' survey in autumn/early winter every year. Some of these surveys are then repeated the following spring as part of a 'Partial Measures' survey.

Each year, an Analytical Report is produced for each individual authority, providing a detailed analysis and interpretation of the 'Full Measures' surveys. This is followed by a brief Update Report for each individual authority, providing ongoing findings from the 'Partial Measures' surveys. Annually, a Cell 1 Overview Report is also produced. This provides a region-wide summary of the main findings relating to trends and interactions along the entire Cell 1 frontage.

To date the following reports have been produced:

Table 1 Analytical, Update and Overview Reports Produced to Date

Year		Full Measures		Partial Measures		Cell 1 Overview Report
		Survey	Analytical Report	Survey	Update Report	
1	2008/09	Sep-Dec 08	May 09	Mar-May 09		-
2	2009/10	Sep-Dec 09	Mar 10	Feb-Mar 10	July 10	-
3	2010/11	Aug-Nov 10	Feb 11	Feb-Apr 11	Aug 1	Sep 11
4	2011/12	Sep 2011	Aug 12	Mar-May 12	Feb 13	
5	2012/13	Sept 2012	Feb 13	Mar-Apr 13	May 2013	
6	2013/14	Oct 2013	Feb 14	Mar-Apr 14	July 14	
7	2014/15	Nov 2014	Feb 15	March 15	June 15	
8	2015/16	Nov 2015	Feb 16	April 16	July 16	
9	2016/17	Aug / Sept 2016	Jan 17 (*)			

(*) The present report is **Analytical Report 9** and provides an analysis of the 2016 Full Measures survey for County Durham Council's frontage.

In addition, separate reports are produced for other elements of the programme as and when specific components are undertaken, such as wave data collection, bathymetric and sea bed sediment data collection, aerial photography, and walk-over visual inspections.

For purposes of analysis, the Cell 1 frontage has been split into the sections listed in Table 2.

Table 2 Sub-divisions of the Cell 1 Coastline

Authority	Zone
Northumberland County Council	Spittal A
	Spittal B
	Goswick Sands
	Holy Island
	Bamburgh
	Beadnell Village
	Beadnell Bay
	Embelton Bay
	Boulmer
	Alnmouth Bay
	High Hauxley and Druridge Bay
	Lynemouth Bay
	Newbiggin Bay
	Cambois Bay
Blyth South Beach	
North Tyneside Council	Whitley Sands
	Cullercoats Bay
	Tynemouth Long Sands
	King Edward's Bay
South Tyneside Council	Littehaven Beach
	Herd Sands
	Trow Quarry (incl. Frenchman's Bay)
	Marsden Bay
Sunderland Council	Whitburn Bay
	Harbour and Docks
	Hendon to Ryhope (incl. Halliwell Banks)
Durham County Council	Featherbed Rocks
	Seaham
	Blast Beach
	Hawthorn Hive
	Blackhall Colliery
Hartlepool Borough Council	North Sands
	Headland
	Middleton
	Hartlepool Bay
Redcar & Cleveland Borough Council	Coatham Sands
	Redcar Sands
	Marske Sands
	Saltburn Sands
	Cattersty Sands (Skinningrove)
Scarborough Borough Council	Staithes
	Runswick Bay
	Sandsend Beach, Uppang Beach and Whitby Sands
	Robin Hood's Bay
	Scarborough North Bay
	Scarborough South Bay
	Cayton Bay
	Filey Bay

1. Introduction

1.1 Study Area

Durham County Council's frontage extends from Ryhope Dene to Crimdon Beck. For the purposes of this report and for consistency with previous reporting, it has been sub-divided into five areas, namely:

- Featherbed Rocks
- Seaham (Dawdon)
- Blast Beach
- Hawthorn Hive
- Blackhall Colliery

1.2 Methodology

Along Durham County Council's frontage, the following surveying is undertaken:

- Full Measures survey annually (since 2008) each autumn/early winter comprising:
 - Beach profile surveys along nine transect lines
- Partial Measures survey annually (since 2009) each spring comprising:
 - Beach profile surveys along six transect lines
- Cliff top survey bi-annually at:
 - Seaham (Dawdon)

The location of these surveys is shown in Figure 2. The 2016 Full Measures survey was undertaken along the Seaham and Easington frontage on the 25th August 2016 and along the Blackhall frontage on the 30th September 2016. During the Seaham & Easington survey the weather was overcast with light rain, with a smooth sea state and a force two wind from the east. During the Blackhall survey the weather was showery and overcast, with a rough sea state and a force three wind from the south west.


All data have been captured in a manner commensurate with the principles of the Environment Agency's *National Standard Contract and Specification for Surveying Services* and stored in a file format compatible with the software systems being used for the data analysis, namely SANDS and ArcGIS. This data collection approach and file format is comparable to that being used on other regional coastal monitoring programmes, such as in the South East and South West of England.

Upon receipt of the data from the survey team, they are quality assured and then uploaded onto the programme's website for storage and availability to others and also input to SANDS and GIS for subsequent analysis.

The Analytical Report is then produced following a standard structure for each authority. This involves:

- description of the changes observed since the previous survey and an interpretation of the drivers of these changes (Section 2);
- documentation of any problems encountered during surveying or uncertainties inherent in the analysis (Section 3);
- recommendations for 'fine-tuning' the programme to enhance its outputs (Section 4); and
- providing key conclusions and highlighting any areas of concern (Section 5).

Data from the present survey are presented in a processed form in the Appendices.



Key

SURVEY LOCATIONS

Topographic Profiles

- Annual
- Bi-Annual

Topographic Surveys

- 6 monthly
- yearly
- 5 yearly

Cliff Top Monitoring Pegs

- @ 300

(Indicative Survey Extents shown)

Client: North East Coastal Group

Project: Cell 1 Regional Coastal Monitoring Programme

Figure 2 - Map 1

Durham County Council Frontage

Analytical Report Topo Surveys


Drawing Scale at A4 1:25,000

WATER
 Royal HaskoningDHV
 Marlborough House
 Marlborough Crescent
 Newcastle upon Tyne
 NE1 4EE

Tel: +44 (0)191 211 1300
 Fax: +44 (0)191 211 1313
 www.royalhaskoningdhv.com



Contains OS data © Crown Copyright and database right 2016



Key

SURVEY LOCATIONS

Topographic Profiles

— Annual

— Bi-Annual

(Indicative Survey Extents shown)

Client: North East Coastal Group

Project: Cell 1 Regional Coastal Monitoring Programme

Figure 2 - Map 2

Durham County Council

Council Frontage

Analytical Report

Topo Surveys

Drawing Scale at A4 1:25,000

WATER

Royal HaskoningDHV

Marlborough House

Marlborough Crescent

Newcastle upon Tyne

NE1 4EE

Tel: +44 (0)191 211 1300

Fax: +44 (0)191 211 1313

www.royalhaskoningdhv.com



Contains OS data © Crown Copyright and database right 2016

2. Analysis of Survey Data

2.1 Featherbed Rocks

Survey Date	Description of Changes Since Last Survey	Interpretation
30 th September 2016	<p>Beach Profiles:</p> <p>One beach profile line 1bEA1, located at Featherbed Rocks (Appendix A), has been monitored since March 2009. The profile extends across the cliff top and cliff face then extends across the promenade (chainage 55), rock armour sea defence (chainage 55 to 80) and beach. At the base of the sea wall rock armour extends as far as 80m chainage. Beyond 80m there has been little change over the summer of 2016, the beach profiles reflect the rocky nature of the foreshore and that there is no beach over the shore platform. Previous surveys have shown accumulations of material at the base of the revetment but this has not been present since the 2012 Full Measures survey.</p>	<p>The rocky nature of this foreshore means it is unlikely to undergo significant changes in morphology unless sediment is deposited upon it. A veneer beach has previously been present here but has not been recorded since the 2012 Full Measures survey.</p> <p>Longer term trends: Between 2010 and 12 a thin veneer beach was present. Since 2013 the profiles recorded have all been low exposing the rocky shore platform along much of its length.</p>

2.2 Seaham (Dawdon)

Survey Date	Description of Changes Since Last Survey	Interpretation
<p>30th September 2016</p>	<p>Cliff-top Survey:</p> <p>Three ground control points have been established along the cliff top at Dawdon (Figure B1). The separation between any two points is nominally 300m. These cliff top surveys are intended to inform on erosion rates of the undefended sea cliffs extending south of the rock armour revetment to the south of Seaham Harbour.</p> <p>The cliff top surveys at Dawdon are undertaken bi-annually. Measurements are taken from a fixed ground control point along a fixed bearing to the edge of the cliff top. Appendix B provides information about the ground control points and results from between the 2008 (baseline) cliff top survey and the current (September 2016) survey.</p> <p>Between April 2016 and September 2016 the most northerly post showed little change, with the middle post showing small amount of recession of 0.1m. The most southerly post showed the largest change with 0.2m of recession.</p> <p>Appendix C provides results from the September 2016 survey, showing the distance from the ground control point to the edge of the cliff top along the defined bearing and changes in position since the November 2008 baseline survey.</p>	<p>Two of the three monitoring locations showed retreat during summer 2016 indicating the cliffs have been locally active.</p> <p>Longer term trends: Long-term recession rates calculated from the data collected since November 2008 show retreat at 0.1m/yr. for Point 1 and 0.2m/yr. at Point 3 at the margins of the bay and no change at Point 2 in the centre of the bay.</p>

2.3 Blast Beach

Survey Date	Description of Changes Since Last Survey	Interpretation
30 th September 2016	<p>Beach Profiles:</p> <p>Blast Beach is covered by four beach profile lines (Appendix A). All of the profiles along Blast Beach exhibit similar forms, with a rock cliff, wide colliery spoil beach with a distinct low cliff at its eroding seaward edge, and a mixed gravel and sand foreshore extending to MLW.</p> <p>Profile 1bSH1b was added to the programme during the Full Measures survey in October 2015. The profile is adjacent to the sewage works south of Seaham. The profile is cliff to 30m and then gravel beach between 30m and 60m chainage, which has shown some slight fluctuation in level over the last year but no long term change. There are two concrete blocks which have been upturned on the beach and are shown on the profiles as a protrusion in the profile between 60m and 65m chainage. The beach is visible again between 65m and 70m chainage. Below this point the rocks are exposed from 70m chainage to the end of the survey at 85m.</p> <p>Profile 1bSH1a was added to the programme during the Full Measures survey in September 2009. It is located to the north of the previously-established 1bSH1. The upper beach has a very similar profile to the previous year as far as the eroding face of the spoil deposit at 140m chainage. Between 140m chainage and 150m there has been little change in the beach profile since April 2016. From 150m to 160m chainage the beach levels have dropped by 1.5m and the crest in the beach has moved landward by 3m. Between 160m and 190m chainage the beach gradient has remained similar but the level has dropped by 1m over the summer of 2016. From 190m chainage and the end of the survey at 260m chainage the rocks are exposed at the bottom of the beach. The 2016 profile is towards the lower end of the range of previously recorded profiles, and is around 1 m lower than the autumn 2015 profile.</p> <p>Profile 1bSH1 is similar to all of the previous surveys to the beach crest at 75m. The mound present at the top of the beach in March 2015 between chainages 90m and 100m has reappeared and moved landwards by 10m. Between 90m and 145m the beach gradient has remained similar since October 2015 however the beach level has dropped by up to 1.2m (0.5m since April 2016). From 145m to the end of the survey at 160m chainage the rocks are exposed. The 2016 profile is towards the lower end of the range of previously recorded profiles.</p>	<p>The cliffs behind Blast Beach are currently inactive because they are fronted by colliery spoil. The crest of the spoil material on profiles 1bSH1 and SH1a has remained stable since 2009. Profile 1bSH2 has been progressively eroding since 2009, however has remained more stable since November 2014 showing accretion in the lower beach.</p> <p>The beaches at profiles 1bSH1b, 1bSH1a, and SH1 have all been subject to up to a 1.5m drop in beach level since the April 2016 survey. The beach at profile SH2 has shown an increase in level however. There has also been some change in the mid beach levels with formation of berms.</p> <p>Longer term trends: The sea cliffs will eventually reactivate as on-going erosion of the colliery spoil removes the protection it affords to the cliffs. This is most likely to occur at the southern end of the bay where the spoil is most rapidly eroding. The accumulating sediment seaward of the colliery spoil in the northern part of the bay will offer the cliffs more protection. However since the winter of 2014 there has been a reversal in the trend with erosion in the north of the bay and accretion in the south; this may yet be a short term change.</p>

Survey Date	Description of Changes Since Last Survey	Interpretation
	<p>Profile 1bSH2 is largely similar to the previous surveys as far as the current beach crest at 120m chainage. The crest in the beach has shown progressive erosion since 2009, with the crest retreating by around 20m. The beach from 135m to 190m chainage has increased by 1m with the development of a slight berm at 150m. At the bottom of the profile at 190m to 200m chainage rocks are exposed on the beach. The 2016 profile is in the middle of the range of previously recorded profiles.</p>	

2.4 Hawthorne Hive

Survey Date	Description of Changes Since Last Survey	Interpretation
30 th September 2016	<p>Beach Profiles:</p> <p>Hawthorne Hive is monitored by beach profile 1cEA2 (Appendix A). The survey report notes "<i>unable to measure start of Section EA2 as the vegetation has choked out the section line and route over cliff faces</i>" and therefore all surveys following October 2012 start at 95m chainage.</p> <p>In previous years there was a channel which crossed the profile; however since April 2013 it has been infilled. The majority of the beach, particularly between 110m and 145m chainage shows an increase in beach levels since the April 2016 profiles, with the berm at chainage 118m having moved seawards slightly by 5m. The rest of the profile between 145m and 220m chainage has the rocks exposed at the bottom of the beach. The beach as a whole is 0.3m above the October 2015 profile.</p>	<p>The beach has recovered since the lowest levels recorded in April and November 2014, and is near the middle of the range of historical levels recorded since 2008.</p> <p>Longer term trends: The beach level has recovered since the lows of 2014 and shows continued increases in levels. Limited cliff erosion occurs in this section and therefore sediment supply is limited to erosion of colliery spoil. Storm events which may block the channel and varying flows in Hawthorne Burn are likely to continue to episodically block the channel and change its course across the beach.</p>

2.5 Blackhall Colliery

Survey Date	Description of Changes Since Last Survey	Interpretation
25 th August 2016	<p>Beach Profiles:</p> <p>Blackhall Colliery is covered by three beach profile lines (Appendix A). As at Blast Beach, profiles are dominated by colliery spoil and exhibit similar forms with a rock cliff, wide spoil beach with a distinct cliff at the eroding face of the colliery spoil, and a gravel and sand foreshore that extends to MLW.</p> <p>1cBH1 is located near Horden Point and shows that the face of the colliery spoil has receded by a similar amount as in 2015 of around 0.5, much less than the 3m erosion, which happened in 2014. The eroding face at 140m chainage is steep. The overall gradient of the beach has changed becoming steeper, with an increase in beach levels of up to 0.4m from 140m chainage to 150m. From 155m to 175m chainage the beach level has dropped by 0.2m since October 2015, making it the lowest recorded levels in this section. From 175m to 200m chainage the cobble beach has remained stable, with an increase in sand levels from 200m to the end of the profile.</p> <p>Profile 1cBH2 exhibits no change in the cliff profile and the cliffed-edge of the spoil beach has remained stable since November 2014. There had been 2m of erosion since September 2013, but much less through 2015 and 2016. There is around c.43m of material from the eroding face at the back of the beach to the cliff toe. From 170m to 250m the beach gradient has reduced, with erosion of up to 0.4m between chainage 170m to 205m, and infilling of the profile by up to 1m between chainage 205m and 250m making the lower beach the highest recorded levels. At the bottom of the profile from chainage 250m the beach has remained stable.</p> <p>The profile 1cBH3 shows that since 2008 there has been episodic migration, infilling and scouring of the outflow of Castle Eden Burn, which crosses the profile. There has been limited recession of the landward bank of the channel, which is closer to the cliffs than in previous years. From the edge of the channel at 150m to 185m chainage there is a mound where the upper beach berm is. The top of the mound has moved landward by 5m. From 180m to 225m there has been erosion of up to 0.5m since October 2015. The lower beach is the lowest recorded level because of the progressive recession of this beach.</p>	<p>The more northerly two profiles at Blackhall Colliery show a similar trend with little change in the eroding face of the beach material following a large change in 2014.</p> <p>There have been some changes to the gradients of the profiles, with BH1 steepening whilst BH2 has flattened slightly, and BH3 has remained consistent.</p> <p>Longer term trends: The surveys show that the spoil beach along much of the Blackhall Colliery shore is progressively eroding but continues to protect the cliffs in the short term.</p>

3. Problems Encountered and Uncertainty in Analysis

The cliff top position surveys at Dawdon are assumed to have a limit of accuracy of $\pm 0.1\text{m}$ due to the techniques used. The accuracy of short-term recession data are therefore limited, but longer-term recession rates will become more reliable as further data is obtained (see section 1.3).

At Blast Beach SH1 and SH1A there was no access to the cliff top due to dense vegetation. At Hawthorne Hive the surveyor was unable to measure the start of Section EA2 as the vegetation has choked out the section line and route over cliff faces. At Blackhall the surveyor was unable to access part of sections BH1 and BH2 due to dense vegetation.

4. Recommendations for 'Fine-tuning' the Monitoring Programme

No changes are recommended at the present time.

5. Conclusions and Areas of Concern

- At Featherbed Rocks the rocky shore platform continues to be exposed and the veneer beach present in earlier surveys has been absent since autumn 2012.
- At Seaham cliffs there has been recession along ground control points 1 and 3 at the margins of the bay of between 0.1m/yr. and 0.2m/yr. since the records began in November 2008. No significant change has occurred at ground control point 2 at the centre of the bay. Further years of data collection will help to understand the long term trends on these cliffs and the stability of the bay.
- At the Blast Beach colliery spoil still prevents the sea from acting directly at the natural cliff toe; however it can be expected that the cliffs will reactivate in coming years following erosion of the spoil deposit. Since winter 2014 there has been a reversal in the long term trends with erosion at the northern end of the bay and accretion at the southern end, which makes it more difficult to predict which section of cliff will reactivate first.
- At Hawthorne Hive the levels on the foreshore have recovered since April and November 2014 and are continuing to increase, they are now in the middle of the range of recorded beach levels. However, it is likely that the long term trend of progressive erosion will continue on this profile.
- At Blackhall Colliery, the seaward face of the colliery spoil deposit continues to erode in the northern part of the bay. In the southern of the bay, mound of beach material continues to erode and the channel has been moving landward. The channel is likely to scour the beach sediments under high flows, but become infilled again by wave action under storm conditions.

Appendices

Appendix A
Beach Profiles

Beach Profile

Location: 1bEA1

Date: 30/09/2016

Inspector: AG

Low Tide:

Low Tide Time:

Wind


Sea State:

Visibility:

Rain:

Summary: 2016 Full Measures Topo Survey

Easting: 442861.92 Northing: 549874.593 Profile Bearing: 50 ° from North



Beach Profile

Location: 1bSH1B

Date: 30/09/2016

Inspector: AG

Low Tide:

Low Tide Time:

Wind


Sea State:

Visibility:

Rain:

Summary: 2016 Full Measures Topo Survey

Easting: 443599.944 Northing: 548130.378 Profile Bearing: 63 ° from North



Beach Profile

Location: 1bSH1

Date: 30/09/2016

Inspector: AG

Low Tide:

Low Tide Time:

Wind


Sea State:

Visibility:

Rain:

Summary: 2016 Full Measures Topo Survey

Easting: 443613.742 Northing: 547404.589 Profile Bearing: 74 ° from North



Beach Profile

Location: 1bSH2

Date: 30/09/2016

Inspector: AG

Low Tide:

Low Tide Time:

Wind


Sea State:

Visibility:

Rain:

Summary: 2016 Full Measures Topo Survey

Easting: 443806.533 Northing: 546899.552 Profile Bearing: 74 ° from North



Beach Profile

Location: 1cEA2

Date: 30/09/2016

Inspector: AG

Low Tide:

Low Tide Time:

Wind


Sea State:

Visibility:

Rain:

Summary: 2016 Full Measures Topo Survey

Easting: 444101.532 Northing: 545888.48 Profile Bearing: 75 ° from North



Beach Profile

Location: 1cBH1

Date: 25/08/2016 Inspector: AG

Low Tide:

Low Tide Time:


Wind Sea State:

Visibility:

Rain:

Summary: 2016 Full Measures Topo Survey

Easting: 444443.313 Northing: 542826.089 Profile Bearing: 71 ° from North



Beach Profile

Location: 1cBH2

Date: 25/08/2016 Inspector: AG

Low Tide:

Low Tide Time:


Wind Sea State:

Visibility:

Rain:

Summary: 2016 Full Measures Topo Survey

Easting: 445046.836 Northing: 541386.805 Profile Bearing: 58 ° from North



Beach Profile

Location: 1cBH3

Date: 25/08/2016

Inspector: AG

Low Tide:

Low Tide Time:

Wind


Sea State:

Visibility:


Rain:

Summary: 2016 Full Measures Topo Survey

Easting: 445771.315 Northing: 540371.473 Profile Bearing: 49 ° from North



Beach Profiles: 1bEA1




Profiles Envelope
03/12/2008
16/10/2015
10/04/2016
30/09/2016

HAT
MHWS


MLWS

SANDS

Beach Profiles: 1bSH1B



Beach Profiles: 1bSH1A




Profiles Envelope
21/10/2009
16/10/2015
10/04/2016
30/09/2016

HAT
MHWS

MLWS

SANDS

Beach Profiles: 1bSH1




Profiles Envelope
25/11/2008
16/10/2015
10/04/2016
30/09/2016

HAT
MHWS

MLWS

SANDS

Beach Profiles: 1bSH2




- Profiles Envelope
- 25/11/2008
- 16/10/2015
- 10/04/2016
- 30/09/2016

HAT
MHWS

MLWS

Beach Profiles: 1cEA2



Profiles Envelope
03/12/2008
16/10/2015
10/04/2016
30/09/2016

HAT
MHWS

MLWS

SANDS

Beach Profiles: 1cBH1




Profiles Envelope
04/11/2008
22/11/2014
15/10/2015
25/08/2016

HAT
MHWS

MLWS

SANDS

Beach Profiles: 1cBH2




Profiles Envelope
04/11/2008
22/11/2014
15/10/2015
25/08/2016

HAT
MHWS

MLWS

SANDS

Beach Profiles: 1cBH3



Profiles Envelope
04/11/2008
22/11/2014
15/10/2015
25/08/2016

HAT
MHWS

MLWS

SANDS

Appendix B
Cliff Top Survey

Cliff Top Survey

Seaham

Three ground control points have been established on the Seaham frontage (Figure B1). The maximum separation between any two points is nominally 300m.

The cliff top surveys at Seaham are undertaken biannually. Measurements are taken from a fixed ground control point along a fixed bearing to the edge of the cliff top.

Table B1 provides baseline information about these ground control points and results from the 2008 (baseline) survey showing the position from the ground control point to the edge of the cliff top along the defined bearing. Future reports will show results from subsequent surveys and provide a means of assessing erosion since the baseline survey.

Table B1 – Cliff Top Surveys at Seaham

Ground Control Points				Distance to Cliff Top (m)			Total Erosion (m)		Erosion Rate (m/year)
Ref	Easting	Northing	Bearing	Baseline Survey	Previous Survey	Present Survey	Baseline to Present	Previous to Present	Baseline to Present
			(°)	Nov 2008	Apr 2016	Sep 2016	Nov 2008 - Sep 2016	Apr 2016 - Sep 2016	Nov 2008 - Sep 2016
1	443515.4	548421.7	70	16.1	15.09	15.05	1.05	0.04	0.13
2	443607.8	548136.3	90	13.3	13.37	13.25	0.05	0.12	0.01
3	443756.1	547858.5	95	14.8	13.54	13.34	1.46	0.20	0.18

INSERT FIG B1